



SCHOOL OF NEW RESOURCES -- IMMUNIZATION RECORD FORM

THIS FORM MUST BE COMPLETED AND RETURNED PRIOR TO REGISTRATION

Name: _____ Date of Birth: _____

Address: _____
(Street) (City) (State) (Zip)

Social Security Number: _____ Phone: _____

E-mail Address: _____

Campus: _____

NEW YORK STATE, PUBLIC HEALTH LAW, CHAPTER 2165 requires the following information regarding MMR immunizations and dates. First dose must be on or after first birthday. Doses must be at least 28 days apart.

MMR (Measles, Mumps, Rubella – Combined) Vaccine

or

Measles (Rubeola) Immunity: Check & complete all that apply:

[] Two doses of live measles vaccine on or after January 1, 1968:

OR

[] Date of immune measles titer:

OR

[] Date of physician diagnosed measles disease:

Form fields for MMR dates: / / (mm/dd/yy) and / / (mm/dd/yy) for two different rows.

and

Mumps Immunity: Check & complete all that apply:

[] One dose of mumps vaccine on or after January 1, 1969:

OR

[] Date of immune mumps titer:

OR

[] Date of physician diagnosed mumps disease:

Form fields for Mumps dates: / / (mm/dd/yy) and / / (mm/dd/yy) for two different rows.

and

Rubella (German Measles) Immunity: Check & complete all that apply:

[] One dose of rubella vaccine on or after January 1, 1969:

OR

[] Date of immune rubella titer:

Form fields for Rubella dates: / / (mm/dd/yy) and / / (mm/dd/yy) for two different rows.

Health Care Provider Signature _____

License Number _____

Date (mm/dd/yy) _____

NEW YORK STATE, PUBLIC HEALTH LAW, CHAPTER 2167 requires that all college and university students provide the following information by checking only one box and signing.

- [] had the meningococcal (Menomune™) vaccine within the past 10 years (revaccinate every 3-5 years if increased risk continues). Date ___/___/___ OR
[] had the meningococcal (Menactra™) vaccine, preferred . Date ___/___/___ OR
[] I have read, or have had explained to me, the information regarding meningococcal meningitis disease. I understand the risks of not receiving the vaccine. I have decided that I (my child) will not obtain immunization against meningococcal meningitis disease.

Student Signature _____

Date (mm/dd/yyyy) _____

PLEASE SUBMIT THIS FORM TO: HEALTH SERVICES, ANGELA HALL

Meningococcal Vaccines

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 What is meningococcal disease?

Meningococcal disease is a serious bacterial illness. It is a leading cause of bacterial meningitis in children 2 through 18 years old in the United States. Meningitis is an infection of the covering of the brain and the spinal cord.

Meningococcal disease also causes blood infections.

About 1,000–1,200 people get meningococcal disease each year in the U.S. Even when they are treated with antibiotics, 10–15% of these people die. Of those who live, another 11%–19% lose their arms or legs, have problems with their nervous systems, become deaf, or suffer seizures or strokes.

Anyone can get meningococcal disease. But it is most common in infants less than one year of age and people 16–21 years. Children with certain medical conditions, such as lack of a spleen, have an increased risk of getting meningococcal disease. College freshmen living in dorms are also at increased risk.

Meningococcal infections can be treated with drugs such as penicillin. Still, many people who get the disease die from it, and many others are affected for life. This is why preventing the disease through use of meningococcal vaccine is important for people at highest risk.

2 Meningococcal vaccine

There are two kinds of meningococcal vaccine in the U.S.:

- Meningococcal conjugate vaccine (**MCV4**) is the preferred vaccine for people 55 years of age and younger.
- Meningococcal polysaccharide vaccine (**MPSV4**) has been available since the 1970s. It is the only meningococcal vaccine licensed for people older than 55.

Both vaccines can prevent 4 types of meningococcal disease, including 2 of the 3 types most common in the United States and a type that causes epidemics in Africa. There are other types of meningococcal disease; the vaccines do not protect against these.

3 Who should get meningococcal vaccine and when?

Routine vaccination

Two doses of MCV4 are recommended for adolescents 11 through 18 years of age: the first dose at 11 or 12 years of age, with a booster dose at age 16.

Adolescents in this age group with HIV infection should get three doses: 2 doses 2 months apart at 11 or 12 years, plus a booster at age 16.

If the first dose (or series) is given between 13 and 15 years of age, the booster should be given between 16 and 18. If the first dose (or series) is given after the 16th birthday, a booster is not needed.

Other people at increased risk

- College freshmen living in dormitories.
- Laboratory personnel who are routinely exposed to meningococcal bacteria.
- U.S. military recruits.
- Anyone traveling to, or living in, a part of the world where meningococcal disease is common, such as parts of Africa.
- Anyone who has a damaged spleen, or whose spleen has been removed.
- Anyone who has persistent complement component deficiency (an immune system disorder).
- People who might have been exposed to meningitis during an outbreak.

Children between 9 and 23 months of age, and anyone else with certain medical conditions need 2 doses for adequate protection. Ask your doctor about the number and timing of doses, and the need for booster doses.

MCV4 is the preferred vaccine for people in these groups who are 9 months through 55 years of age. MPSV4 can be used for adults older than 55.



4**Some people should not get meningococcal vaccine or should wait.**

- Anyone who has ever had a severe (life-threatening) allergic reaction to a previous dose of MCV4 or MPSV4 vaccine should not get another dose of either vaccine.
- Anyone who has a severe (life threatening) allergy to any vaccine component should not get the vaccine. *Tell your doctor if you have any severe allergies.*
- Anyone who is moderately or severely ill at the time the shot is scheduled should probably wait until they recover. Ask your doctor. People with a mild illness can usually get the vaccine.
- Meningococcal vaccines may be given to pregnant women. MCV4 is a fairly new vaccine and has not been studied in pregnant women as much as MPSV4 has. It should be used only if clearly needed. The manufacturers of MCV4 maintain pregnancy registries for women who are vaccinated while pregnant.

Except for children with sickle cell disease or without a working spleen, meningococcal vaccines may be given at the same time as other vaccines.

5**What are the risks from meningococcal vaccines?**

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of meningococcal vaccine causing serious harm, or death, is extremely small.

Brief fainting spells and related symptoms (such as jerking or seizure-like movements) can follow a vaccination. They happen most often with adolescents, and they can result in falls and injuries.

Sitting or lying down for about 15 minutes after getting the shot—especially if you feel faint—can help prevent these injuries.

Mild problems

As many as half the people who get meningococcal vaccines have mild side effects, such as redness or pain where the shot was given.

If these problems occur, they usually last for 1 or 2 days. They are more common after MCV4 than after MPSV4.

A small percentage of people who receive the vaccine develop a mild fever.

Severe problems

Serious allergic reactions, within a few minutes to a few hours of the shot, are very rare.

6**What if there is a serious reaction?****What should I look for?**

Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS is only for reporting reactions. They do not give medical advice.

7**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

8**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

**Vaccine Information Statement (Interim)
Meningococcal Vaccine**

10/14/2011

42 U.S.C. § 300aa-26

Office Use Only

